



High Harmonic spectroscopy unveils the phase transitions of high-temperature superconductors

Researchers from ICFO, ICMAB-CSIC and Guangdong Technion-Israel Institute of Technology have developed a new methodology to investigate and measure the quantum phase transitions of a high-temperature superconductor by using High Harmonic spectroscopy.

September 27, 2022

Superconductors are materials that exhibit the ability to conduct electricity without any resistance. This phenomenon is observed in materials when they are cooled below the so-called superconductor transition temperature, often at very low temperatures (a few degrees above the absolute 0). Among these materials, there are the so-called high-temperature superconductors, which behave as superconductors at temperatures above 77K (the boiling point of liquid nitrogen). These materials are showing to be essential in the development of new electronic and information processing devices as well as optical

quantum computers and even for improving the efficiency of electrical transmission lines. However, high-temperature superconductivity has been seen to be closely linked to the control of their microscopic dynamics. So far, the detection of the different microscopic quantum phases in these complex materials has resulted quite challenging. Not only are the physical processes of these dynamic states still incomplete due to their wide array of quantum states, but the current methods used to explore their dynamics at microscopic scales are lagging sensitivity. Therefore, new tools to better understand the dynamic evolution of these types of superconductors are needed.

Now, in an international study, ICFO researchers Utso Bhattacharya, Ugaitz Elu, Tobias Grass, Piotr T. Grochowski, Themistoklis Sidiropoulos, Tobias Steinle, and Igor Tyulnev, led by ICREA Professors Jens Biegert and Maciej Lewenstein, in collaboration with ICMA-B-CSIC researchers Jordi Alcala and Anna Palau, and Marcelo Ciappina, from the Guangdong Technion-Israel Institute of Technology, propose a new methodology based on the use of High Harmonic spectroscopy (HHS) to investigate the transitions between the different phases of YBCO, copper oxide cuprate material which is a well-known high-temperature superconductor. This study represents a major scientific breakthrough since it is the first time that high non-linear and non-perturbative diagnostics/detection methodology is used to understand the behavior of strongly correlated materials

In view of the experimental results obtained, the researchers have also gone beyond and present a new theoretical model to identify the connection between the measured optical spectra and the transition between the different quantum states of the YBCO: strange metal pseudogap, and superconductor. The study has been recently published in the journal PNAS. In their experiment, the researchers used 100nm thick films of YBCO mounted on a micro-refrigerator. Firstly, they characterized the superconducting properties of the YBCO films and confirmed their quality. Then, by using ultra-short infrared laser pulses the author of the study induced high harmonics generation in the material samples, which were placed inside a vacuum chamber and cooled to a temperature of 77K

High Harmonics are the high-energy photons emitted by the electrons of a system when it is placed in a strong laser field. These emitted photons have a frequency many times that of the driving laser field

When they hit the surface, they recorded the reflected radiation with a spectrograph to study the harmonic spectrum, which contains the imprints of this nonlinear optical response, and it was found to have a connection with the phase transitions

Seeing these experimental results in the lab and the lack of a theory that could explain what was being observed, the researchers developed a new strong-field quasi-Hubbard model that shed light on the connections between the measured high harmonics and the formation of Cooper pairs, that is, the paired electrons that are responsible for the superconducting phase

When using this new theoretical model, the theoretical calculations of the high harmonics

spectra obtained matched the experimental data. "The model faithfully reproduces the functional form of the measurement data over the entire temperature range and for several orders of magnitude of harmonic amplitude", the authors highlighted. This new approach, as they noted, has permitted a theoretical connection between the measurements and the underlying microscope dynamics providing a "powerful new methodology to study the quantum phase transitions" in correlated materials.

Finally, the team highlights that their work provides a "first striking example" of how High Harmonic Spectroscopy can be used to distinguish correlated phases of matter. They also consider that it paves the way toward a "refined understanding of the physical processes that occur inside high-temperature superconductors".

Acknowledgements

We thank V. Kunets from MMR Technologies and X. Menino at Institut de Ciències Fòniques for technical support and Dr. K. Dewhurst and Dr. S. Sharma for help with the ELK code. J.B., U.E., T.P.H.S., T.S., and I.T. acknowledge financial support from the European Research Council (ERC) for ERC Advanced Grant *TRANSFORMER* (grant 788218) and ERC Proof of Concept Grant *miniXi* (grant 840010). J.B. and group acknowledge support from FET-OPEN *PETACOMI* (grant 829153), FET-OPEN *OPTOLOGIC* (grant 899794), EIC-2021-PATHFINDEROPEN *TwistedNano* (grant 101046424), Laerlab-Europe (grant 654148), Marie Skłodowska-Curie ITN *smart-Xi* (grant 860553), Plan Nacional PID-PID2020-112664GB-I00-210901, AGAUR for 2017 SGR 1639, *Severo Ochoa* (grant SEV-2015-0522), Fundacio Cellex Barcelona, the CECA Programme/Generalitat de Catalunya, and the Alexander von Humboldt Foundation for the Friedrich Wilhelm Bessel Prize. U.B., T.G., P.T.G., and M.L. acknowledge funding from the ERC for ERC Advanced Grant NOQIA; Agencia Estatal de Investigación (R&D project CEX2019-000910-S, funded by MCIN/AEI/10.13039/501100011033, Plan Nacional FIDEU PID2019-106901GB-I00, FPI, QUANTERA MAQS PCI2019-111828-2, Proyectos de Investigación *Retos Colaboración* RTC2019-007196-7); Fundacio Cellex; Fundacio Mir-Puig; Generalitat de Catalunya through the CERCA program; AGAUR grant 2017 SGR 134; QuantumCAT U16-011424, cofunded by ERDF Operational Program of Catalonia grant 201-2020; EU Horizon 2020 FET-OPEN OPTOLOGIC (grant 899794); National Science Centre, Poland (Symfonia grant 2016/20/W/ST4/00314); Marie Skłodowska-Curie grant S REDCH 101029393; *La Caixa* Junior Leaders fellowships (ID10001044); and EU Horizon 2020 under Marie Skłodowska-Curie grant agreement 847648 (LCF/BQ/119/11690013, LCF/BQ/PI20/11760031, LCF/BQ/PR20/11770012, and LCF/BQ/PR21/1184013). P.T.G. acknowledges the Polish National Science Center grants 2018/31/N/ST2/1429 and 2020/36/T/ST2/00065 and is supported by the Foundation for Polish Science. Center for Theoretical Physics of the Polish Academy of Sciences is a member of the National

Laboratory of Atomic, Molecular and Optical Physics. A.P. and J.A. acknowledge support from the Spanish Ministry of Economy and Competitiveness through the Severo Ochoa Programme for Centres of Excellence in R&D (grant SEV-2015-0496), SuMaTe project RTI2018-095853-B-C21) cofinanced by the European Regional Development Fund, and the Catalan Government grant 2017-SGR-1519. This work was supported by EU COST action NANOCOHBRI CA16218. M.C. acknowledges financial support from the Guangdong Province Science and Technology Major Project (Future functional materials under extreme conditions - 2021B0301). We also acknowledge the Scientific Services at Institut de Ciència de Materials de Barcelona and ICN2.