

Felicidades al nuevo graduado de doctorado del ICFO

El Dr. Arturo Villegas se ha graduado con una tesis titulada ?Optical parameter sensing: sensitivity limits and the advantages of using spatial modes of light'

April 26, 2023

Felicitamos el Dr. Arturo Villegas que hoy ha defendido su tesis en la sala Homage del ICFO. El Dr. Villegas obtuvo su master en Fisica Aplicada por la Delft University of Technology, en los Paises Bajos. Se unio como estudiante de doctorado en el grupo de investigacion de Quantum Engineering of Light en ICFO dirigido por el Profesor Juan P Torres. La tesis del Dr. Villegas titulada ?Optical parameter sensing: sensitivity limits and the advantages of using spatial modes of light' fue supervisada por el Profesor Juan Perez Torres.

RESUMEN:

Resolution enhancement in modern optical metrology techniques has been possible due to significant technological improvements. The accurate and precise control of wavelength,

bandwidth, and power of light sources; homogeneity, high-quality composition materials and surface smoothness of optical elements; and highly stable nanopositioners and optomechanical components, allow more sensitive imaging and sensing, in certain scenarios even beyond the standard diffraction limit. This has motivated a more detailed study of the fundamental resolution limitations of an optical system.

The chosen approach to address this problem is to consider optical imaging as a parameter estimation problem. With this in mind, the theory of quantum estimation and statistical inference provides the tools to determine the estimation precision limits. Starting by considering the state of light as a quantum state that carries information of interest; the Cramer-Rao lower bound provides a fundamental limit for the achievable precision. This lower bound is directly associated to optical resolution in practical terms.

In this thesis, we present an overview of the useful tools of quantum estimation theory that can be applied to optical metrology. We focus on the Cramer-Rao lower bound, and provide methods to calculate it. Since the bound depends on specific characteristics of the system, we explore three specific possibilities.

First we present with an example the validity regimes for different bounds; explicitly to calculate the parameters of entangled photon pairs in the frame of a quantum Lidar System. Second, we show the dependency of the lower bound on the photonic model selection, showing a discrepancy between N copies of a single photon and a multimode coherent state with average photon number N.

two particular models. Third, we study the effects of lossy environments in the informational content of the quantum state.

Additionally, we present the conditions that a measuring strategy must satisfy to allow attainability of the fundamental limit.

If the measurement strategy is not feasible or it is not possible to implement, one can evaluate the resolution improvement of the available technique by comparing to the standard and fundamental limits.

For specific scenarios of interest, measurement methods based on the use of spatial modes of light allow to asymptotically attain the resolution limit. This has drawn attention to the information carried by specific modes, and has motivated the design of measurement strategies based on probing or sensing using spatial modes. In this thesis we include an overview of spatial modes of light, their generation and detection, and their use for optical sensing. We present a method for optical beam localization in the transverse plane using spatial mode information. Moreover, we propose a technique to retrieve the full modal decomposition of an arbitrary beam; which combined with the adequate set of modes allows to estimate certain parameters of an optical state with the maximum precision possible.

Comite de Tesis:

Dr. Pablo Loza, ICFO

Dr. Etienne Brasselet, Laboratoire Ondes et Matiere d'Aquitaine (LOMA)

Dr. Armand Eugene Albert Koolen, ASML Research



Comite de Tesis: